

## **Chapter IV: The Wildlife Habitat Response Model: A Tool for Estimating Terrestrial Wildlife Habitat Responses to Fuel Treatments**

Authors: David S. Pilliod,<sup>1,2</sup> Katharine Shick,<sup>3</sup> M. Elena Velasquez<sup>4</sup>

<sup>1</sup>Aldo Leopold Wilderness Research Institute, Rocky Mountain Research Station, USDA Forest Service, P.O. Box 8089, Missoula, Montana 59807

<sup>2</sup>Department of Biological Sciences, College of Science and Mathematics, California Polytechnic State University, San Luis Obispo, CA 93407; Phone: (805) 756-2207; Email: dpilliod@calpoly.edu

<sup>3</sup>927 Cooper Street, Missoula, MT 59802

<sup>4</sup>3008 S Longleaf, Boise, ID 83716

Manuscript will be published as a chapter in the following General Technical Report: Pilliod, D.S., K. Shick, and M.E. Velasquez. *In review*. The Wildlife Habitat Response Model: a tool for estimating terrestrial wildlife habitat responses to fuel treatments. Pages xx-xx in Sutherland, E.K., and A.E. Black (eds). Estimating the environmental consequences of fuel treatments: user's guide. USDA Forest Service General Technical Report GTR-RMRS-xx, Rocky Mountain Research Station, Missoula, MT.

Model Location:

<http://forest.moscowfsl.wsu.edu/fuels/tools.html>

**Abstract**

This paper describes the Wildlife Habitat Response Model (WHRM), a web-based computer tool for evaluating the potential effects of fuel reduction projects on terrestrial wildlife habitats in dry coniferous forests of the western United States. WHRM uses species-habitat associations to predict how fuel treatments may affect species habitats. Using data gleaned from the scientific literature, WHRM first identifies the habitat elements, such as down wood or snags, which are important for species in terms of the following life history requirements: reproduction, food acquisition, and shelter from predators and environmental extremes. Managers then enter the proportional amount of change from pretreatment conditions for each habitat element listed. Decisions about change amounts can be derived from fuel treatment objectives, desired future conditions, or predictions from forest stand development models. WHRM then calculates a weighted average effect of each specified treatment for each life history category based on the changes in individual habitat elements weighted by literature support. The output of WHRM is strictly qualitative (for example, will management activity result in a positive, negative, or have no effect on habitat for a species). The model output can help managers compare alternative treatments and quickly identify potential negative effects of treatments on wildlife habitats. This information, along with an annotated bibliography, provides a starting point for developing the environmental consequences components of NEPA-type assessments for fuel treatment activities.

## **Introduction**

Managers face a difficult task in predicting the effects of fuel treatments on wildlife within the dry interior forests of the western United States. Few empirical studies are available from which to draw inferences and thus there is uncertainty and some concern about how sensitive wildlife species will respond to mechanical thinning, prescribed burning and their alternatives (Pilliod and others 2006). Anecdotal observations suggest that wildlife mortality during forest thinning and prescribed burning operations is minimal and inconsequential to populations of terrestrial species (Folk and Bales 1982; Komarek 1969). Populations are more likely to respond to the rapid changes and sometimes prolonged recovery in forest structure and composition that result from fuel management activities. When empirical studies on the effects of fuel treatments on wildlife habitats are unavailable, some predictions may still be possible by first identifying the habitat requirements of a species and then estimating how fuel management activities will alter the habitat elements that are important to a species' survival and reproduction. The Wildlife Habitat Response Model (WHRM) is a web-based predictive computer tool that was developed to meet this need.

WHRM is based on species-habitat relationships and organized similarly to an envirogram. Species-habitat relationships are descriptive mechanisms used by wildlife biologists and managers for linking species with the habitats in which they are associated. Typically organized in matrices and databases, species-habitat relationships can be general (for example, species-biome) or specific (for example, species-habitat element) and can have qualifiers on habitat usage such as resident, seasonal, occasional, breeding, foraging, and other descriptive terms. An envirogram is a graphic representation of causal relationships linking indirect to direct causes of species responses to the environment (Andrewartha and Birch 1984). Each causal pathway is listed under one of four categories: mates, resources, predators, and mal-entities. These four categories represent the general habitat requirements of a species and can be described as those features or habitat elements that are needed for an animal to (1) find a mate, reproduce, and successfully rear offspring (for example, breeding sites, birthing areas, and nest sites), (2) acquire the nutrition necessary for survival and reproduction (for example, foraging habitat, forage, and habitat for prey which influences prey availability), (3) escape predation, and (4) seek shelter from environmental hazards (for example, daytime or nighttime temperatures, extreme weather events, seasonal climate fluctuations, and unpredictable disturbances such as drought, fire, or flooding).

The goal of this paper is to provide a user's guide to the Wildlife Habitat Response Model. In the following sections we describe the concept, inputs, outputs, utility, limitations, and assumptions of the model. In chapter XI of this document, we provide a case-study of how WHRM can be used for fuel treatment planning.

## **Description**

The goal of our modeling effort was to produce a straightforward tool to help fuel planners and NEPA specialists (that is, non-biologists) to conceptualize and qualitatively predict how fuel reduction treatments might affect the habitats of terrestrial wildlife species living in dry, coniferous forest ecosystems of the western U.S. Hence, WHRM is intended for planning purposes and general assessments and is not intended to replace detailed population-level assessments of threatened, endangered, or sensitive species.

WHRM was designed to predict potential habitat changes relative to pre-treatment conditions and can be used to compare the potential effects of different treatments on a species' habitats under each of the following life history requirements: reproduction, food resources, and shelter.

In essence, WHRM packages a wildlife-habitat database into a user-friendly computer-based tool. WHRM first identifies the habitat requirements of a chosen species and then allows users to investigate how proposed fuel management activities and alternatives may influence the critical habitat needs of a species. The predicted response of a species to proposed habitat changes is based on species-habitat associations reported in the scientific literature for ponderosa pine (*Pinus ponderosa*) and dry-type Douglas-fir (*Pseudotsuga menseizii*), lodgepole pine (*Pinus contorta*) and mixed conifer forests in the western U.S. (see map and definition, pages xx-xx, this document). Therefore, WHRM predictions are specific to these forest types and may not apply to other forest types or a species across its entire range. Furthermore, WHRM predictions are most appropriate within the stand being modeled and should not be generalized to landscape levels. Predictions have no temporal scale, but users can produce time-specific predictions based on the values selected for changes or recovery in each habitat element. Predictions from forest stand development models, such as the Forest Vegetation Simulator with the Fire and Fuels Extension (FVS-FFE) can aid in predicting how various habitat elements may change over time.

Species-habitat associations used in this model were generated from an extensive search of over 450 peer-reviewed published articles on wildlife and wildlife habitats. In most cases, we used original data from papers published in scientific journals. For each paper, we recorded the location of the study, habitats investigated, whether the paper was a disturbance paper (in other words, describing habitat associations in a disturbed or recovering environment), species studied, the response variable used (for example, occurrence, abundance), a description of the habitat elements with which a species was significantly correlated (probability less than 0.05 or 0.10, depending on the usage in the paper), and the direction of the correlation. Habitat elements were then placed into standard categories (see table 1). If the habitat categories described in a paper and those used in WHRM did not match exactly, we used a liberal, inclusive approach. For example, if a paper reported snags in the size class 12 to 24 inches diameter-at-breast-height (d.b.h.), we used size classes 10 to 19 inches and 20 to 29 inches d.b.h. for WHRM. If size classes were not provided in a paper, we used a general category (for instance, snags (size not specified)). Based on information reported in a paper, we placed each habitat association described for a species into one of four categories: reproduction, non-consumptive foraging habitat, forage or prey habitat, and shelter from predators or environmental extremes. Some habitat associations fell into two or more categories depending on the life history of the animal. If not specified in a paper, we categorized remaining habitat associations based on field guides, species accounts, and other general references. Data from papers outside the target area were not included in WHRM unless no other information was available. Under such circumstances, we first included papers from similar forest types (for example, moist, coastal Douglas-fir) within the region and then from other forest types (for example, spruce).

## Mechanics

### Inputs

#### *Step 1: Select fuel treatments to compare*

The first step to running WHRM is to select two to four fuel treatments to compare from the following list: Thinning and Pile Burning, Thinning and Broadcast Burning, Prescribed Fire with No Thinning, and Wildfire. Wildfire is considered the no-action alternative.

#### *Step 2: Select a species*

Only one species can be modeled at a time. WHRM provides a drop-down list of species occurring in dry interior forest types of the western U.S. for which there is sufficient life history information to predict responses to habitat alterations.

#### *Step 3: Create input tables*

After selecting the fuel treatment type and the species, click on the button *Display Habitat Associations*. WHRM creates an input table for each treatment selected and displays the habitat elements (see definitions in table 1) that are considered important habitat requirements for the life history of the selected species (figure 1). Some habitat elements will be listed under more than one life history requirement category.

#### *Step 4: Indicate change in habitat elements*

WHRM requires users to input how each fuel treatment will alter the specific habitat elements that are associated with a species. The user specifies the magnitude and direction of change in each habitat element based on one or a combination of the following sources: (1) fuel treatment objectives, (2) desired future conditions, or (3) simulated results using computer models such as FVS-FFE with the WHRM compute variable (see appendix A) and the Understory Response Model for shrubs, forbs, and grasses (Sutherland and Miller, this volume). The change in habitat elements is a categorical measure of the proportional change in a habitat element relative to pre-treatment conditions (figure 1).

The above approach works well for habitat elements that are reasonably common in a stand, but requires careful consideration for habitat elements that are rare. If a habitat element is at low in abundance in a stand, it will show an exaggerated proportional response with small increases or decreases in abundance. For example, if there are only 2 trees in the 20 to 29 cm d.b.h. size class in your stand and one of them dies during a burn, then that mortality will result in a 50% decrease in abundance of that size class of trees. This may or may not be biologically meaningful, depending on the species of interest. Therefore, users need to be aware of the absolute abundance of various habitat elements and make careful decisions when determining the proportional change in habitat elements that are rare in a stand. We recommend selecting “no change” for habitat elements that are at very low abundance in the stand, unless these are absolutely critical to a particular wildlife species.

#### *Step 5: Run model*

After the expected change in each habitat element has been selected in each treatment table, click on the button *Run Wildlife Habitat Response Model*. A treatment

comparison table is generated from the user inputs that shows the predicted average weighted effect on habitat for each treatment divided into each life history category (figure 2). These are qualitative predictions and listed as: highly positive, moderately positive, slightly positive, neutral, slightly negative, moderately negative, or highly negative (see table 4). To understand how habitat element changes are influencing these average effects, users should examine the treatment-specific tables by clicking on the treatment title (figure 3).

WHRM generates the habitat element response prediction by multiplying the value given to the relative magnitude of change expected in a given habitat element (table 2) by the expected association of that habitat element with a given species (table 3). For example, in figure 3 under reproduction, WHRM multiplied the values -2 and 1 to generate a value of -2 (which equates with a moderately negative effect, table 4), because mule deer (*Odocoileus hemionus*) are positively associated (value of 1) with shrub cover for fawning habitat and shrub cover decreased by 41-70% (value of -2) in this treatment. Predicted habitat changes are then averaged within life history requirements weighted by the “confidence” in the habitat association. “Confidence” was quantified as the number of studies demonstrating a given association. Therefore, habitat elements that are found to be important in several studies are given more weight than habitat elements that are only cited once. Weighted average habitat changes were placed into descriptive categories (table 4).

Averaging across each life history requirement provides an estimate of change in habitat for reproduction, forage, or cover, but an average may not always be a meaningful summary and we recommend looking at how the change in each habitat element influences a species’ habitat and discuss these with a wildlife biologist. To examine how each habitat element is contributing to an average, users can view each treatment table separately by clicking on the treatment name in the output table (figure 3).

There will be cases where, within one life history requirement (for example, foraging habitat), the change in some habitat elements results in a positive influence on habitat suitability whereas the change in other habitat elements results in a negative influence. When this occurs, WHRM flags the output with a cautionary statement suggesting that users carefully evaluate the individual habitat elements and/or discuss them with a locally knowledgeable biologist to decide which habitat elements are most important given the existing conditions within the stand and surrounding landscape.

## Outputs

The output of WHRM is a table showing the average weighted habitat effects for each specified treatment subdivided into life history requirements for reproduction, foraging, and cover. Figure 3 shows an example of an output table. In this treatment, the removal of smaller trees and subsequent increase in grass and forbs resulted in increased forage for mule deer despite the negative effects of shrub removal for cover.

The output table is followed by a summary statement. The summary statement first identifies those studies that were used to build the habitat associations. Users can find information about these studies following the summary statement. The summaries describe the information gleaned from each publication that we considered relevant to WHRM and may not match the overall conclusions of the paper or even original

objectives. The table and text in each output can be cut and pasted into a planning document.

The summary statement also identifies those treatments that resulted in a negative effect on a species' reproduction, foraging, or cover habitats. To aid the user, the output summary also identifies those habitat elements that contributed to these negative habitat changes. Finally, we provide a table of general information for each species, including: distribution, habitat preferences, diet, predators, home range sizes, responses to disturbance, general sources of information used, specific sources of information used in the model, and summaries of selected scientific publications.

### **Interpretation and Assumptions**

WHRM is most appropriately used to assist managers in decisions about how fuel treatments can be applied such that the most important habitat elements for wildlife species of interest are protected or enhanced. By organizing species-habitat associations by reproduction, food acquisition, and shelter, WHRM enables users to see what part of an animal's life history may be most affected by a fuel treatment. The information in WHRM output also can be used by wildlife biologists to identify information gaps and help direct future research. As is the case with all models, the quality of the prediction is only as good as the data available to run the model and information on some species is limited. As new or revised information on species-habitat associations becomes available, the simple structure of WHRM programming should enable easy and frequent updates.

The primary assumption behind WHRM is that habitat associations reported in the literature are correct (Table 5). One of the potential problems with such an assumption is that most correlative information reported in field studies is untested. Correlative studies generally use an exploratory hindcasting approach to explain patterns observed in species occurrence and abundance at a specific place and period of time. Given that most of these correlations have never been validated with novel datasets, there is certain risk in using them to make general predictions about species responses to habitat change because environmental, demographic, and ecological conditions may vary considerably among locations or over time (see additional discussion in Morrison and others 1992). Although we cannot account for these variable or unmeasured factors, the user can reduce the uncertainty of their predictions by only applying WHRM to make predictions in appropriate forest types and by checking the background information used to build the model for each species.

The second assumption of WHRM is that habitat associations are linear and essentially one to one. In other words, a 11-40% increase in a positively associated habitat element results in a 11-40% increase in habitat suitability. Assumptions that all responses are linear can be misleading. For instance, some species may prefer intermediate levels of canopy closure, avoiding closed canopy and open forests. We did not address non-linear relationships in this version of the model. If a species selected intermediate levels of a habitat element, we generally did not include it in the models. Instead, this information was described under "General Habitat Associations" in the Background Information output.

The third assumption of WHRM is that species abundance or probability of occurrence is correlated with habitat suitability. For the most part, the habitat associations in WHRM are based on a species' abundance in relation to a habitat element.

However, high abundance does not necessarily mean high quality habitat, especially when considering altered habitats (Van Horne 1983). Estimates of vital rate parameters (for instance, fecundity, survival, dispersal) are needed to make a true assessment of habitat quality or potential source and sink habitats. However, these more complicated modeling procedures are only possible with fairly extensive field data on local populations and thus are generally beyond the scope of most fuel plans and NEPA assessments.

### **Limitations**

The accuracy of the WHRM predictions will vary depending on the species examined. WHRM will likely work best for species that are closely associated with forest structure and composition or have very specific habitat needs (for example, trees of a specific diameter for nesting). WHRM may provide meaningful information for some but not all habitats of species that use very different habitats seasonally (for example, elk), use multiple habitats for different life history traits (for example, lynx denning in continuous mature forest but hunting along edges), or are extremely far-ranging (for example, grizzly bears). Habitat generalists and species not closely associated with forest structure and composition will likely not respond to fuel treatments (Pilliod and others, 2006) and in such cases, WHRM may not provide much useful information. There are several factors that can influence predictions of habitat suitability. The following paragraphs discuss these factors and limitations to WHRM.

The reliability of WHRM predictions may be strongly influenced by climate patterns such as drought (Fulé and others 2002), post-treatment weather such as intense rainstorms (Robichaud, unpublished data), forest type, landscape physiognomy, subtleties of fuel treatment activities, and other factors. Overall, our confidence in suggested patterns decreases with time since treatment. For example, shrubs are likely to decrease considerably (for example, more than 40%) 1 year after broadcast burning, but then fully recover within 5-10 years depending on the type shrub and level of mortality (Sutherland and Miller, this volume). However, this assumes shrubs are top-killed only and are free from herbivory. Shrubs requiring seed regeneration will take longer and herbivory from large ungulates, like mule deer and Rocky Mountain elk (*Cervus elaphus*), can slow recovery rates of understory shrubs following thinning and/or prescribed burning (Huffman and Moore 2003, 2004).

Another limitation to consider is the scale or extent of predictions. This issue has two facets. First, with the popularity of landscape-level analysis and ecosystem approaches to wildlife management, it may be tempting to use WHRM for purposes beyond its intent and scope. WHRM is intended to be a stand-level predictive tool and should not be extrapolated to broader spatial scales. Second, many of the species included in WHRM do not use one stand for all of their habitat needs. Therefore, WHRM predictions may be less informative for a species that has a large home range and can simply avoid unfavorable habitat conditions without cost. We draw attention to this issue by comparing the size of the proposed treatment area with the typical home range size for a species and provide this information in the output. Hence, the predicted model results only apply for conditions within the stand being altered. Surrounding stand conditions and larger landscape characteristics may ameliorate or exacerbate the predicted results.



Finally, WHRM does not produce quantitative predictions of population responses to fuel treatments nor does it provide a measure of prediction confidence. If a fuel treatment may pose a significant risk to the habitats of a species of concern, we recommend using more specific modeling approaches such as various habitat suitability models or risk assessment tools such as population viability models (for examples, see Roloff and others 2001). Keep in mind, however, that these more complex models require at least some information on habitat patch location, quantity, and quality, and/or information on population sizes, age structures, and vital rates to estimate parameters that are meaningful. This level of information is rarely available for most species and locations.

Despite these limitations, species-habitat matrices are generally considered favorable relative to other habitat-based wildlife modeling approaches because of their practical application, simple model structure, and generality and communicability of model output (table 1 in Roloff and others 2001). WHRM is intended to be easy to understand and use for non-biologists, and yet provide meaningful general predictions about the potential environmental consequences of fuel treatments. As fuel reduction planning is stream-lined and more projects are rapidly moved through the review process, WHRM provides a tool to help managers quickly assess the potential effects of specific fuel treatments on wildlife habitats.

#### **Acknowledgements**

The programming for this model was developed by Elena Velasquez. Thanks to Stephanie Rebain for providing the coding for the FVS compute variable. We thank The Wildlife Society for coordinating a scientific review and the anonymous reviewers that provided helpful comments on the model and user's guide.

#### **Literature Cited**

Andrewartha, H.G.; Birch, L.C. 1984. *The Ecological Web: More on the Distribution and Abundance of Animals*. Chicago, IL: University of Chicago Press. 506 p.

Folk, R.H. III; Bales, C.W. 1982. An evaluation of wildlife mortality resulting from aerial ignition prescribed burning. *Proceedings Annual Conference Southeastern Association Fish and Wildlife Agencies*. 36: 643-646.

Fulé, P.Z.; Covington, W.W.; Smith, H.B.; Springer, J.D.; Heinlein, T.A.; Huisinga, K.D.; Moore, M.M. 2002. Comparing ecological restoration alternatives: Grand Canyon, Arizona. *Forest Ecology and Management*. 170: 19-41.

Huffman, D.W.; Moore, M.M. 2003. Ungulate herbivory on buckbrush in an Arizona ponderosa pine forest. *Journal of Range Management*. 56: 358-363.

Huffman, D.W.; Moore, M.M. 2004. Responses of Fendler ceanothus to overstory thinning, prescribed fire, and drought in an Arizona ponderosa pine forest. *Forest Ecology and Management*. 198: 105-115.

Komarek, E.E. Sr. 1969. Fire and animal behavior. Proceedings Annual Tall Timbers Fire Ecology Conference. 9: 160-207.

Morrison, M.L.; Marcot, B.G.; Mannan, R.W. 1992. Wildlife-habitat relationships: Concepts and applications. Madison, WI: The University of Wisconsin Press. 343 p.

O'Neil, T.A.; Bettinger, K.A.; Heyden, M.V.; Marcot, B.G.; Barrett, C.; Mellen, T.K.; Vanderhaegen, W.M.; Johnson, D.H.; Doran, P.J.; Wunder, L.; Boula, K.M. 2001. Structural conditions and habitat elements of Oregon and Washington. In: Johnson, D.H.; O'Neil, T.A., eds. Wildlife-habitat relationships in Oregon and Washington. Corvallis, Oregon: Oregon State University Press: 115-139.

Pilliod, D.S.; Bull, E.L.; Hayes, J.L.; Wales, B.C. 2006. Wildlife and invertebrate response to fuel reduction treatments in dry coniferous forests of the western United States: a synthesis. Gen. Tech. Rep. RMRS-GTR-173. Ft. Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 34 p.

Roloff, G.J.; Wilhere, G.F.; Quinn, T.; Kohlmann, S. 2001. An overview of models and their role in wildlife management. In: Johnson, D.H.; O'Neil, T.A., eds. Wildlife-habitat relationships in Oregon and Washington. Corvallis, Oregon: Oregon State University Press: 512-536.

Sutherland, S.; Miller, M. 2005. Predicting the impact of fuels reduction on understory species: the Understory Response Model. In: Sutherland, E.K.; Black, A.E., eds. Estimating the environmental consequences of fuels treatments: user's guide. Gen. Tech. Rep. RMRS-GTR-xx. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: xx-xx.

Van Horne, B. 1983. Density as a misleading indicator of habitat quality. Journal of Wildlife Management. 47: 893-901.

Table 1. List of habitat elements important for wildlife. Definitions of habitat elements adapted from O’Neil and others (2001). Units are not displayed because WHRM uses percent change from pre-treatment conditions and thus any units can be used.

Forest Component	Habitat Elements	Definition
Forest Floor	Bare Mineral Soil Exposure	The inorganic soil layer beneath the humus. This HE is usually expressed as percent exposure in a given area representative of the stand. Although not generally considered a wildlife HE, bare mineral soil represents a lack of duff and litter cover (1-(duff+litter cover)) and therefore is indicative of poor habitat quality for some species and increased erosion potential.
	Duff Cover	The matted layer of organic debris beneath the litter layer. Decomposition is more advanced than in litter layer; intergrades with uppermost humus layer of soil. This HE is usually expressed as percent cover of a given area representative of the stand. Duff depth can also be important for wildlife habitat, but is not included.
Understory Vegetation	Grass Cover	The amount of ground cover composed of grasses. This HE is usually expressed as percent cover of a given area representative of the stand.
	Forb/Herbaceous Cover	The cover of understory, non-woody vegetation layer beneath the shrub layer that includes forbs, mosses, and ferns. This HE is usually expressed as percent cover of a given area representative of the stand.
	Shrub Cover	A measure of shrub density and usually visually estimated as a vertical projection of shrub crown diameter onto the ground. Shrubs are further subdivided into short shrubs (0-18” tall) and tall shrubs (>18” tall). Tall shrubs are usually considered ladder fuels whereas short shrubs are not. If shrub height is not specified in a study, it is listed under "shrub cover (all size classes)". Shrub species and number of shrub canopy layers are important for wildlife habitat, but are not specified.

Forest Component	Habitat Elements	Definition
Down Wood	Litter Cover	The upper layer of loose, organic (primarily vegetative) debris on the forest floor. Decomposition may have begun, but components still recognizable. This HE is usually expressed as percent cover of a given area representative of the stand. WHRM does not specify litter depth, but this can influence wildlife use.
	Down Wood	Includes downed logs, branches, and rootwads. This HE is usually expressed in tons per acre and described as downed woody debris (DWD). DWD is subdivided in size classes based on maximum diameter: 0-3" (considered fine woody debris, FWD and includes 1, 10 and 100 hour fuels) and >3" (considered coarse woody debris, CWD and includes all 1000 hour fuels). WHRM subdivides CWD in size classes 4-6", 7-12", and >12". If a study does not specify size classes, then it is listed under "down wood (all size classes). The WHRM does not specify decay class, but this can be important for wildlife use.
Standing Dead Wood	Snags	Standing dead wood can be grouped (listed as "snags (all size classes)") or separated into size classes measured in diameter at breast height (dbh) or at 4.5 feet above the ground. WHRM groups snags into the following size classes: 0-4.9" dbh, 5-9" dbh, 10-19" dbh, 20-29" dbh, and 30+" dbh. Snags are usually expressed in snags per acre. Stumps can be considered short snags. WHRM does not specify decay class, but this can be important for wildlife use.
Live Trees	Crown Base Height	The height from the ground to the base of the live overstory tree canopy. This HE is usually expressed as a linear measure, such as feet or meters.
	Tree Canopy Cover	The cover of the coniferous and deciduous living tree canopy projected vertically to the ground. This HE is usually expressed as a percentage.
	Trees	Living trees are usually described as a density (trees per acre or TPA). All coniferous trees are grouped together (listed as "trees (all species and size classes)") or grouped by size classes measured as dbh (see snags). WHRM also lists aspen trees and "other trees", which are usually site specific such as Gambel Oaks.

Table 2. Ordinal values used to quantify the change in habitat elements associated with fuel treatments.

Change in Habitat Element	Value Used in Computation
Decrease more than 70%	-3
Decrease 41-70%	-2
Decrease 11-40%	-1
No change (+/- 10%)	0
Increase 11-40%	1
Increase 41-70%	2
Increase more than 70%	3

Table 3. Values used to quantify the association of a species with a habitat element. Relationships were estimated from correlative studies with response variables including probabilities of occurrence, indices of abundance (number observed or densities), and population estimation.

Relationship between a species and a habitat element	Value Used in Computation
Negative	-1
Neutral	0
Positive	1

Table 4. Values used to depict species habitat responses to planned fuel treatments.

Value Range	Description
-3.0 to -2.1	Highly Negative
-2.0 to -1.1	Moderately Negative
-1.0 to -0.26	Slightly Negative
-0.25 to 0.25	Minimal to None
0.26 to 1.0	Slightly Positive
1.1 to 2.0	Moderately Positive
2.1 to 3.0	Highly Positive

Table 5. Underlying assumptions in the Wildlife Habitat Response Model.

Assumptions of the Wildlife Habitat Response Model

- 
1. Habitat associations reported in the literature are correct.
  2. Habitat associations are linear and mostly one to one.
  3. Abundance or probability of occurrence are correlated with habitat suitability
-



Figure 1. Example of habitat element input fields. User decides the relative change in each habitat element based on treatment objectives, desired future conditions, or values derived from computer simulation models.

Figure 2. Example of the output generated by the Wildlife Habitat Response Model. The output is read left to right. In this case, two treatments are compared: Thinning and Broadcast Burning and Thinning and Pile Burning.

Figure 3. By clicking on the treatment title *Thinning and Broadcast Burning* in Figure 2, users can view the treatment-specific table that was used to calculate the *weighted average effect on habitat* for each life history category for each treatment. Users will need to refer to these treatment-specific tables to determine how habitat element changes are influencing average effects, particularly when positive habitat element changes and negative habitat element changes are neutralizing average effects and increasing uncertainty.

Fig 1.



**Wildlife Habitat Response Model**



**Habitat Associations for  
Mule deer (*Odocoileus hemionus*)**

Select the change in each habitat element based on expected results of each planned fuel treatment.  
Each fuel treatment has its own input table.

**Thinning and Broadcast Burning**

Life History Requirements		Key Wildlife Habitat Elements	Change in Habitat Elements						
			greater than 70% Decrease	41-70% Decrease	11-40% Decrease	No Change (+ / - 10%)	11-40% Increase	41-70% Increase	greater than 70% Increase
Reproduction	Nest Sites, Birthing Areas, Breeding Sites	Forb/Herbaceous Cover	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>
		Shrub Cover 0-18" height	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
		Trees (all species) 0-4.9" dbh	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Food Resources	Foraging Habitat	Tree Canopy Cover	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
	Forage, Prey Habitat	Grass Cover	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
		Forb/Herbaceous Cover	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>
		Shrub Cover (size classes not specified)	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Cover	Shelter from Predators and Environmental Extremes	Shrub Cover (size classes not specified)	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
		Tree Canopy Cover	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
		Shrub Cover 18+" height	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Fig. 2



**Wildlife Habitat Response Model**  
**Comparison of Selected Fuel Treatments**



Click on fuel treatment titles to view details about specific predictions for habitat elements and life history requirements

**Taxonomic group:** Ungulates  
**Species:** Mule deer (*Odocoileus hemionus*)

Life History Requirements	Key Habitat Element(s)	Weighted Average Effect on Habitat	
		Thinning and Broadcast Burning	Thinning and Pile Burning
Nest Sites, Birthing Areas, Breeding Sites	Forb/Herbaceous Cover	Moderately Negative, but low confidence *	Slightly Negative
	Shrub Cover 0-18" height		
	Trees (all species) 0-4.9" dbh		
Foraging Habitat	Tree Canopy Cover	Moderately Negative	Moderately Negative
Forage, Prey Habitat	Grass Cover	Slightly Positive, but low confidence *	Slightly Positive
	Forb/Herbaceous Cover		
	Shrub Cover (size classes not specified)		
Shelter from Predators, and Environmental Extremes	Shrub Cover (size classes not specified)	Moderately Negative	Slightly Negative
	Tree Canopy Cover		
	Shrub Cover 18+'' height		

\*When the "weighted average predicted effect on habitat" includes some habitat elements that positively influenced habitat and others that negatively influenced habitat, then the average may be misleading. Due to this uncertainty, we recommend you consult a wildlife biologist regarding the relative importance of each habitat element contributing to averages that have low confidence.

Fig 3



**Wildlife Habitat Response Model**  
**Thinning and Broadcast Burning Details**



Click back arrow to return to model comparison

Taxonomic group:	Ungulates
Species:	Mule deer ( <i>Odocoileus hemionus</i> )

Life History Requirements		Key Habitat Element (s)	Change in Habitat Element (s) (user's selection)	Predicted Effects on Habitat ?	Weighted Average Effect on Habitat ?
Reproduction	Nest Sites, Birthing Areas, Breeding Sites	Forb/Herbaceous Cover	11-40% Increase	Slightly Positive	Moderately Negative, but low confidence *
		Shrub Cover 0-18" height	41-70% Decrease	Moderately Negative	
		Trees (all species) 0-4.9" dbh	Greater than 70% Decrease	Highly Negative	
Food Resources	Foraging Habitat	Tree Canopy Cover	41-70% Decrease	Moderately Negative	Moderately Negative
		Grass Cover	41-70% Increase	Moderately Positive	Slightly Positive, but low confidence *
		Forb/Herbaceous Cover	11-40% Increase	Slightly Positive	
Cover	Shelter from Predators, and Environmental Extremes	Shrub Cover (size classes not specified)	41-70% Decrease	Moderately Negative	Moderately Negative
		Tree Canopy Cover	41-70% Decrease	Moderately Negative	
		Shrub Cover 18+ " height	Greater than 70% Decrease	Highly Negative	

\* When the "weighted average predicted effect on habitat" includes some habitat elements that positively influenced habitat and others that negatively influenced habitat, then the average may be misleading. Due to this uncertainty, we recommend you consult a wildlife biologist regarding the relative importance of each habitat element contributing to averages that have low confidence.  
Click back arrow to return to model comparison

Documentation and User Manual: [Use guide (14-page PDF)] [Exclude (1-page PDF)] [Complete WHRM Bibliography (39-page PDF)]  
The Wildlife Habitat Response Model: WHRM  
Input Interface v. 2005.02.23 (for review only) by Elena Velazquez & David Hall  
Model developed by: David Pilling  
Team leader: Elaine Kennedy Sutherland, USDA Forest Service, Rocky Mountain Research Station



Appendix A. Instructions for using FVS-FFE to generate WHRM input data in the proper format.

1. Set the time scale for the simulation to be at least 12 years with 1 year cycles. So if the inventory date is 2004, you should run the simulation from 2004 to 2016 with one year cycles. In this case, 2004 would be for your initial conditions. The treatments would be done in 2005. The 2006 values would be 1-year post treatment. The 2010 values would be 5 years post treatment. And the 2015 values would be 10 years post treatment. If your inventory date is 1990, you can still do the simulation from 2004 to 2016. FVS will grow the trees up to 2004. Normally, it is not recommended to run FVS on 1-year cycles, but acceptable for 12 years, but not much longer.
2. To get the desired variables, the WHRMCompVar.kcp file needs to be inserted into the run. This is done by going to the Edit Simulation window. You insert it into a simulation by clicking on "Insert from file" and selecting WHRMCompVar.kcp.
3. Make sure the Fire and Fuels Extension to FVS is actually used. This simply means making sure that an FFE keyword, such as FuelOut, is included in the run.
4. If any treatments are being simulated, add the appropriate keywords.
5. Lastly, you need to add the Compute 1 post processor into the simulation.