


| | | |
|-----------|-----------|--------|
| Apo | Syal h.t. | |
| | Phma h.t. | |
| P.sme | Caru h.t. | Caru |
| | Phma h.t. | Phma |
| | Smst | |
| Aogr | Phma h.t. | Phma |
| | Libo h.t. | Libo |
| | Asca h.t. | Asca 1 |
| | Tabr | Tabr |
| | Mefe | |
| | Setr h.t. | |
| | Phma | |
| | Clun h.t. | Clun |
| | Tabr | |
| | Mefe | |
| Xete h.t. | Vagl | |
| Cooc | | |
| Ind | Clun h.t. | Clun 1 |
| | Tabr | |
| | Mefe | |
| | Xete | |
| | Asca h.t. | Asca 2 |
| | Tabr | 1 |
| | Mefe | 1 |
| Gydr h.t. | 4 | |
| Adpe h.t. | 10 | |
| Atfi h.t. | Adpe 10 | |
| Atfi | 3 | |
| Opho h.t. | 4 | |
| Tsh | Clun h.t. | Clun + |
| | Arnu | |
| | Mefe | |
| | Xete | |
| | Asca h.t. | Asca |
| Arnu | | |
| Mefe | | |
| Gydr h.t. | | |
| Tsm | Clun h.t. | Mefe |
| | Xete | |
| | Mefe h.t. | Xete |
| | Luh | |
| | Stam h.t. | Mefe |
| Luh | | |
| Xete h.t. | Vagl | |
| Luh | | |
| Abia | Clun h.t. | Clun |
| | Mefe | Mefe |
| | Xete | Xete |
| | Cooc | |
| | Vasc h.t. | Vasc |
| | Luh | |
| | Stam h.t. | Mefe |
| Lica | | |
| Caca h.t. | Caca | |
| Vagl | | |
| Legl | | |
| Cooc | | |
| Xete h.t. | Vagl | |
| Vasc | | |
| Luh | | |
| Pico | Vasc h.t. | |
| | Xete c.t. | |
| | Vaca c.t. | |



POLYPODIACEAE
Adiantum pedatum L.
Adpe

maidenhair fern

FORM: Erect, perennial fern, 10-60cm (4-24in) tall, from short, creeping rhizomes.

LEAVES: Few to solitary deciduous fronds, the stalk divides into two recurving parts, each part consisting of 5-6 pinnae with 15-35 alternate, bluish-green pinnules (leaflets) in a fanlike blade arrangement (1), more or less parallel to ground on a thin, shiny purple to purplish-black wiry petiole.

SORI: Borne on reflexed margins (2).

NOTES: Major populations located between the St. Joe and Selway River drainages, in warm, high humidity habitats.