


Pico	Syal h.t.	
Pico	Phma h.t.	
Pisme	Caru h.t.	Caru
	Aruv	
	Phma h.t.	Phma
	Sms	
Abgr	Phma h.t.	Phma
	Cooc	
	Libo h.t.	Libo
	Xete	
	Asca h.t.	Asca
	Tabr	
	Mefe	
	Setr h.t.	
	Phma	
	Clun h.t.	Clun
	Tabr	
	Mefe	
	Xete	
Xete h.t.	Vagl	
	Cooc	
Thpl	Clun h.t.	Clun
		Tabr
		Mefe
		Xete
	Asca h.t.	Asca
		Tabr
		Mefe
	Gydr h.t.	
	Adpe h.t.	
	Atfi h.t.	Adpe
	Atfi	
Opho h.t.		
	+	
Tshe	Clun h.t.	Clun
		Arnu
		Mefe
		Xete
	Asca h.t.	Asca
	Arnu	
	Mefe	
Gydr h.t.		
Tsmc	Clun h.t.	Mefe
		+
	Xete	
	Mefe h.t.	Xete
	Stam h.t.	Luhi
	Xete h.t.	Vagl
	Luhi	
Abja	Clun h.t.	Clun
		Mefe
		Xete
		Cooc
	Mefe h.t.	Xete
		Vasc
		Luhi
	Stam h.t.	Mefe
		1
		Lica
	1	
Caca h.t.	Caca	
	Vaca	
	Legl	
	Cooc	
Xete h.t.	Vagl	
	Vasc	
	Luhi	
Pico	Vasc h.t.	
	Xete c.t.	
	Vaca c.t.	



LYCOPODIACEAE  
*Lycopodium annotinum* L.  
Lyan

stiff club-moss

FORM: Prostrate, perennial, evergreen clubmoss.

STEMS: Erect, once or twice branched, leafy stems, 5-25cm (2-10in) tall; trailing stems, sparsely leaved, elongate up to 1m (3ft) long, rooting at intervals.

LEAVES: Ascending to lax, crowded in 8 ranks; 4 leaves in each pseudo-whorl, lance-shaped, slightly toothed with very sharp pointed tip, 5-11mm long.

CONES: Sessile, 1-4cm (<2in) long, borne singly at end of some stems.

SORI: None; sporangia reniform, borne on sporophylls, special straw-colored to greenish leaves different from the vegetative leaves, 3-4mm long.

SIMILAR SPECIES: *Lycopodium clavatum* (Stag's horn moss) and other *Lycopodium* species