

Pico	Syal h.t.	1
Pico	Phma h.t.	
Pisme	Caru h.t.	Caru 1
	Phma h.t.	Phma 1
		Smst 1
Abgr	Phma h.t.	Phma 3
		Cooc 3
	Libo h.t.	Libo 1
		Xete 1
	Asca h.t.	Asca 5
		Tabr 3
		Mefe 1
	Setr h.t.	4
		Phma 5
	Clun h.t.	Clun 2
	Tabr 1	
	Mefe 1	
	Xete 4	
Xete h.t.	Vagl 3	
	Cooc 3	
Thp	Clun h.t.	Clun 4
		Tabr 1
		Mefe 1
		Xete 2
	Asca h.t.	Asca 4
		Tabr 2
		Mefe 3
	Gydr h.t.	1
	Adpe h.t.	4
	Atfi h.t.	Adpe 4
	Atfi 4	
Opho h.t.	1	
Tshe	Clun h.t.	Clun 2
		Arnu 2
		Mefe 3
		Xete 3
	Asca h.t.	Asca 4
	Arnu 1	
	Mefe 1	
Gydr h.t.	3	
Tisme	Clun h.t.	Mefe 2
		Xete 4
	Mefe h.t.	Xete 1
		Luhi 2
	Stam h.t.	Mefe 1
		Luhi 1
Xete h.t.	Vagl 1	
	Luhi 1	
Abia	Clun h.t.	Clun 1
		Mefe 1
		Xete 3
	Mefe h.t.	Cooc 3
		Xete 1
		Vasc 1
		Luhi 1
	Stam h.t.	Mefe 2
		Lica 1
	Caca h.t.	Caca 2
	Vaca 1	
	Legl 1	
	Cooc 1	
Xete h.t.	Vagl 1	
	Vasc 1	
	Luhi 1	
Pico	Vasc h.t.	
	Xete c.t.	
	Vaca c.t.	

P t a q



POLYPODIACEAE

Pteridium aquilinum (L.) Kuhn in von der Decken
Ptaq

bracken fern

FORM: Medium to large, erect fern, 5-20dm (2-7ft) tall; vigorously clonal, from deep, branched rhizomes having two light-colored external stripes and distinctive vascular bundles in cross-section.

LEAVES: Deciduous but with a firm, leatherlike feel, tripinnate blade 3-10dm (1-3ft) long with alternate, sessile pinnules, edges rolled under and hairy below, petiole erect, hairy and nonscaly, generally shorter than blade.

SORI: Protected by narrow inrolled indusial leaf margins (1).

ENEMIES: Fire: resistant, regeneration from rhizomes within 1-2 years to preburn levels.

NOTES: Allelopathic to other plant species, and a very aggressive invader of disturbed sites. Reported to contain carcinogenic compounds and is poisonous to livestock.