



POLYPODIACEAE
Woodsia oregana D.C.
Woor

woodsia, Oregon woodsia

FORM: Erect, small to medium tufted fern, 7-25cm (3-10in) tall, from short creeping rhizomes covered with yellowish to dark reddish-brown scales.

LEAVES: Numerous, deciduous, bi to tripinnate, lanceolate blades, 4-15cm (2-6in) long, with 7-17 pairs of opposite or offset pinnae, the lowest one or two pairs smaller than those above, the upper ones gradually reduced to tip (1); persistent petiole bases (2) remain on rhizome.

SORI: Borne on lower surface of the leaf removed from the margins; indusium platelike, with radiating, hairlike often somewhat beaded segments (3), partly concealed by the reflexed margins.

SIMILAR SPECIES: *Cystopteris fragilis*, p.88, without persistent petioles; and *Woodsia scopulina* (Rocky Mtn. woodsia) hairy, glandular leaves.

HABITAT: Talus slopes and cliff crevices and ledges.

	Syal h.t.		Pipo
	Phma h.t.		
	Caru	Caru	Psmc
	Aruv	h.t.	
	Phma	Phma	
	Smst	h.t.	
	Phma	Phma	Abgr
	Cooc	h.t.	
	Libo	Libo	
	Xete	h.t.	
T	Asca	Asca	
+	Tabr		
i	Mefe		
	Setr	h.t.	
	Phma		
T	Clun	Clun	
	Tabr		
i	Mefe		
	Xete		
	Vagl		
	Cooc	Xete	
		h.t.	
T	Clun	Clun	Thp1
	Tabr		
	Mefe		
	Xete		
T	Asca	Asca	
	Tabr		
T	Mefe		
	Gydr	h.t.	
	Adpe	h.t.	
	Adpe	Atfi	
	Atfi	h.t.	
	Opho	h.t.	
	Clun	Clun	Tshe
	Arnu		
	Mefe		
	Xete		
	Asca		
	Arnu		
	Mefe		
	Gydr	h.t.	
	Mefe	Clun	Tsmc
	Xete		
	Xete		
	Luhi		
	Mefe		
	Luhi		
	Vagl	Xete	
	Luhi	h.t.	
	Clun	Clun	Ab1a
	Mefe		
	Xete		
	Cooc		
	Xete	Mefe	
	Vasc		
	Luhi		
	Mefe	Stam	
	Lica		
	Caca	Caca	
	Vaca		
	Legl		
	Cooc		
	Vagl	Xete	
	Vasc		
	Luhi		
	Vasc	h.t.	Pico
	Xete	c.t.	
	Vaca	c.t.	