

PDO	Syal h.t.
	Phma h.t.
Psme	Caru h.t.
	Aruv
	Phma h.t.
	Smst
Phma h.t.	Phma
	Cooc
Libo h.t.	Libo
	Xete
Asca h.t.	Asca 1
	Tabr 1
	Mefe 1
Abgr	Setr h.t. 8
	Phma
	Clun
Clun h.t.	Tabr
	Mefe
	Xete 2
Thp	Asca h.t. 1
	Tabr 1
	Mefe 6
	Gydr h.t.
	Adpe h.t. 2
	Atfi Adpe
	h.t. Atfi 6
	Opho h.t. 1
Tsha	Clun
Clun h.t.	Arnu
	Mefe
	Xete
	Asca h.t. +
	Arnu 3
	Mefe
	Gydr h.t. 1
Tsing	Clun h.t. 2
	Xete 1
	Mefe Xete
	h.t. Luhi 2
Stam	Mefe 7
h.t.	Luhi 5
Xete	Vagl 1
h.t.	Luhi 1
Clun h.t.	Clun
	Mefe 1
	Xete
Mefe h.t.	Cooc 1
	Xete
	Vasc
	Luhi 1
Stam h.t.	Mefe 6
	Lica 9
Caca h.t.	Caca 8
	Vaca 5
	Legl 4
	Cooc
Xete h.t.	Vagl
	Vasc +
	Luhi 1
Pico	Vasc h.t. .
	Xete c.t.
	Vaca c.t.

Setr



ASTERACEAE (COMPOSITAE)

Senecio triangularis Hook.

Setr

arrowleaf groundsel

FORM: Tall, deciduous, perennial forb, 3-15dm (1-5ft) with several stems arising from fibrous roots.

LEAVES: Numerous, alternate, simple, distinctly triangular leaves, 4-20cm (2-8in); basal leaves strongly toothed, with long petioles; cauline leaves become reduced, sessile and less triangular upwards.

FLOWERS: Few to numerous yellow heads in flat-topped inflorescence; June-Sept.

FRUIT: Achene.

HABITAT: Seasonally wet or very moist to riparian sites.