


Pipo	Syal h.t.	1	
	Phma h.t.	7	
Psime	Caru	Caru	
	h.t.	Aruv	
	Phma	Phma 3	
	h.t.	Smst 7	
Abgr	Phma	Phma 7	
	h.t.	Cooc 9	
	Libo	Libo 6	
	h.t.	Xete 4	
	Asca	Asca 9	
	h.t.	Tabr 8	
		Mefe 9	
	Setr	h.t.	10
	Clun	Phma	7
		h.t.	Clun 7
		Tabr 5	
		Mefe 7	
		Xete 7	
Xete	Vagl 2		
h.t.	Cooc 3		
Thpl	Clun	Clun 6	
	h.t.	Tabr 1	
		Mefe 5	
		Xete 3	
	Asca	Asca 7	
	h.t.	Tabr 7	
		Mefe 10	
	Gydr	h.t.	7
	Adpe	h.t.	8
	Atfi	Adpe 7	
h.t.	Atfi 7		
Opho	h.t.	7	
Tshe	Clun	Clun 4	
	h.t.	Arnu 4	
		Mefe 7	
		Xete 5	
	Asca	Asca 9	
h.t.	Arnu 6		
	Mefe 7		
Gydr	h.t.	6	
Tsmc	Clun	Mefe 6	
	h.t.	Xete 5	
	Mefe	Xete	
	h.t.	Luhi 1	
	Stam	Mefe 5	
	h.t.	Luhi 2	
Xete	Vagl 1		
h.t.	Luhi 1		
Abia	Clun	Clun 7	
	h.t.	Mefe 7	
		Xete 7	
		Cooc 3	
	Mefe	Xete	
	h.t.	Vasc	
		Luhi	
	Stam	Mefe 5	
	h.t.	Lica 8	
		Caca 3	
Caca	Vaca		
h.t.	Legl 1		
	Cooc 2		
Xete	Vagl 1		
h.t.	Vasc +		
	Luhi 1		
Pico	Vasc	h.t.	
	Xete	c.t.	
	Vaca	c.t.	



POACEAE (GRAMINEAE) Festuceae
Bromus vulgaris (Hook.) Shear
Brvu

columbia brome

FORM: Robust, hairy, fibrous-rooted perennial, nonrhizomatous, culms slender 30-80cm (12-32in) tall, never forms bunches.

LEAVES: Drooping, broad, flat blades 5-10mm wide, sheaths closed, usually hairy especially at the throat; auricles lacking; membranous ligules with ragged margins (1), 3-5mm long.

INFLORESCENCE: Narrow, open, often drooping, panicle, 10-15cm (4-6in) long; more than 2 florets per spikelet; lemmas awned (2) 3-8mm long; late June-Aug.

ENEMIES: Fire: very resistant, regeneration from seed to preburn levels within 1-2 years.

SIMILAR SPECIES: *Bromus inermis* (smooth brome), rhizomatous.