

C a c a



POACEAE (GRAMINEAE) Agrostideae
Calamagrostis canadensis (Michx.) Beauv.
 Caca

bluejoint reedgrass

FORM: A robust, cool season perennial from creeping rhizomes, sometimes forming tufts; culms hollow, 5-10dm (2-3ft) tall.

LEAVES: Numerous, flat, blades, dull bluish-green, rather lax, rough to touch, 4-8mm wide, open sheath without hairy collar; auricles lacking; finely ciliolate ligules (1), 3-8mm long.

INFLORESCENCE: Open, diffuse panicle, drooping with age, 10-25cm (4-10in) long; mature spikelets have purplish-red color, one floret per spikelet and several spikelets per panicle branch; straight awn, attached at center of lemma, callus hairs nearly as long as lemma (2); late June-Aug.

HABITAT: Moist, poorly drained, cold sites, generally associated with high elevation wet sites.

Syal h.t.		Ptpo	
Phma h.t.			
Caru	Caru	Psmc	
Aruv	h.t.		
Phma	Phma		
Sms t	h.t.		
Phma	Phma	Abyr	
Cooc	h.t.		
Libo	Libo		
Xete	h.t.		
Asca	Asca		
Tabr			h.t.
Mefe			
Setr	h.t.		
Phma	Clun		
Clun			h.t.
Tabr			
Mefe			
Xete			
Vagl		Xete	
Cooc	h.t.		
Clun	Clun	Thp	
Tabr			h.t.
Mefe			
Xete			
Asca	Asca		
Tabr			h.t.
Mefe			
Gydr	h.t.		
Adpe	h.t.		
Adpe	Atfi		
1 Atfi	h.t.		
1 Opho	h.t.		
+ Clun	Clun	Tshe	
Arnu			h.t.
1 Mefe			
Xete			
Asca	Asca		
Arnu			h.t.
Mefe			
Gydr	h.t.		
Mefe	Clun	Tsmc	
Xete			h.t.
Xete			Mefe
Luhi			h.t.
Mefe	Stam		
5 Luhi	h.t.		
Vagl	Xete		
Luhi	h.t.		
Clun	Clun	Ab1a	
Mefe			h.t.
Xete	Mefe		
1 Cooc			h.t.
Xete			
Vasc			
Luhi	Stam		
1 Mefe			h.t.
1 Lica			
10 Caca	Caca		
10 Vaca		h.t.	
4 Legl			
1 Cooc	Xete		
Vagl		h.t.	
+ Vasc			
Luhi			
Vasc	h.t.	Pico	
Xete	c.t.		
Vaca	c.t.		