


Pipo	Syal h.t.	2	
	Phma h.t.	1	
Psme	Caru h.t.	Caru	
	Aruv		
	Phma h.t.	Phma 1	
	Smst	1	
Abgr	Phma h.t.	Phma	
	Cooc		
	Libo h.t.	Libo	
	Xete	1	
	Asca h.t.	Asca 1	
	Tabr	2	
	Mefe	3	
	Setr h.t.	6	
	Clun h.t.	Phma	
		Clun	+
Tabr		3	
Mefe		2	
Xete		2	
Xete h.t.	Vagl	1	
	Cooc	1	
Thpl	Clun h.t.	Clun	
	Tabr		
	Mefe		
	Xete		
	Asca h.t.	Asca	+
	Tabr	1	
	Mefe		
	Gydr h.t.		
	Adpe h.t.		
	Atfi h.t.	Adpe	
	Atfi		
Opho h.t.			
Tshe	Clun h.t.	Clun	
	Arnu	2	
	Mefe		
	Xete		
	Asca h.t.	Asca	1
Arnu			
Mefe			
Gydr h.t.			
Tsmc	Clun h.t.	Mefe	+
	Xete	1	
	Mefe h.t.	Xete	1
	Luhi	1	
	Stam h.t.	Mefe	
Luhi			
Xete h.t.	Vagl	1	
Luhi	1		
Abja	Clun h.t.	Clun	
	Mefe	1	
	Xete	2	
	Cooc		
	Mefe h.t.	Xete	
	Vasc	1	
	Luhi	1	
	Stam h.t.	Mefe	1
	Lica	1	
	Caca h.t.	Caca	2
Vaca			
Legl			
Xete h.t.	Cooc	1	
	Vagl	2	
	Vasc	1	
	Luhi	1	
Pico	Vasc h.t.	1	
	Xete c.t.		
	Vaca c.t.		



CYPERACEAE
Carex rossii Boott
Caro

Ross sedge

FORM: Grasslike perennial with 3-ranked leaves and slender, solid, triangular culms, 1-3dm (4-12in) tall, densely tuft forming from short ascending rhizomes, culms same length or shorter than leaves.

LEAVES: Blades elongate and lax, 1-4mm wide, often longer than culm, basal sheaths reddish.

INFLORESCENCE: Terminal, solitary, staminate spike (1); 1-4 short, few-flowered lateral pistillate spikes borne near staminate spike, pistillate flowers (2) with 3 stigmas, the lowest subtended by a leafy bract (3) which is longer than the staminate spike, one or more small pedunculate pistillate spikes often hidden near base of culms (4); May-Aug.

FRUIT: 3-sided achene, perigynium (5) finely hairy, 2-5mm long, beak 1-2mm long.

ENEMIES: Fire: resistant, regeneration from rhizomes to preburn levels within 1-2 years.

SIMILAR SPECIES: *Carex Conspectus*, p.202.