


Pipo	Syal h.t.	
	Phma h.t.	
Psme	Caru h.t.	Caru
		Aruv
	Phma h.t.	Phma
		Smst 2
Abgr	Phma h.t.	Phma
		Cooc 1
	Libo h.t.	Libo 1
		Xete
	Asca h.t.	Asca +
		Tabr
		Mefe
	Setr h.t.	1
		Phma 1
		Clun 2
Clun h.t.		Tabr
		Mefe
		Xete +
		Vagl
		Cooc
Thp1		Clun 2
	Clun h.t.	Tabr 2
		Mefe
		Xete 2
		Asca 2
	Asca h.t.	Tabr 1
		Mefe
	Gydr h.t.	1
	Adpe h.t.	1
	Atfi h.t.	Adpe 3
	Atfi 1	
Upho h.t.		
Tshe		Clun
	Clun h.t.	Arnu
		Mefe
		Xete 1
		Asca 1
Asca h.t.	Arnu	
	Mefe	
Gydr h.t.		
Tsmc		Clun
	Clun h.t.	Mefe
		Xete
	Mefe h.t.	Xete
		Luhi
	Stam h.t.	Mefe
	Luhi	
Xete h.t.	Vagl	
	Luhi	
Ab1a		Clun
	Clun h.t.	Mefe 1
		Xete 1
		Cooc
	Mefe h.t.	Xete
		Vasc
		Luhi
	Stam h.t.	Mefe 1
		Lica
	Caca h.t.	Caca
	Vaca	
	Legl	
Xete h.t.		Cooc
		Vagl
		Vasc
	Luhi	
Pico	Vasc h.t.	
	Xete c.t.	
	Vaca c.t.	



M e s u



POACEAE (GRAMINEAE) Festuceae
Melica subulata (Griseb.) Scribn.
Mesu

Alaska oniongrass

FORM: Tufted, perennial grass, 30-80cm (12-32in) tall; hollow culms enlarged and bulbous at base (1), attached to short rhizomes.

LEAVES: Very hairy, flat blades, 2-7mm wide, sheaths closed to the top; auricles lacking; membranous ligules (2), 1-5mm long.

INFLORESCENCE: Narrow panicle 10-20cm (4-8in) long, more than 2 florets per spikelet; lemma awnless, finely hairy near base; May-July.

SIMILAR SPECIES: *Bromus vulgaris*, p.199, lemmas awned; and *Melica bulbosa* (onion grass).