

Pipo	Syal h.t.	
Pipo	Phma h.t.	
Psmc	Caru h.t.	Caru
	Aruv	
Psmc	Phma h.t.	Phma
	Smst	
Phma h.t.	Phma	
	Cooc	
Libo h.t.	Libo	
	Xete	
Asca h.t.	Asca	
	Tabr	
	Mefe	
Abgr	Setr h.t.	1
	Phma	
Clun h.t.	Clun	
	Tabr	
	Mefe	1
	Xete	
Xete h.t.	Vagl	
	Cooc	
Clun h.t.	Clun	
	Tabr	
	Mefe	1
	Xete	
Thp1	Asca h.t.	Asca
	Tabr	
	Mefe	
Gydr h.t.		
Adpe h.t.		
Atfi h.t.	Adpe	
	Atfi	
Opho h.t.		
Clun h.t.	Clun	
	Arnu	
	Mefe	
	Xete	
Ishe	Asca h.t.	Asca
	Arnu	
	Mefe	
Gydr h.t.		
Clun h.t.	Clun	
	Mefe	
	Xete	
	Luh1	1
Isme	Stam h.t.	Mefe 2
	Luh1	
Xete h.t.	Vagl	
	Luh1	
Clun h.t.	Clun	
	Mefe	1
	Xete	
	Cooc	4
Mefe h.t.	Xete	2
	Vasc	
	Luh1	
Ab1a	Stam h.t.	Mefe 1
	Lica	1
	Caca	3
	Vaca	10
Caca h.t.	Legl	10
	Cooc	1
	Vagl	
Xete h.t.	Vasc	1
	Luh1	1
	Vasc h.t.	
Pico	Xete c.t.	
	Vaca c.t.	



ERICACEAE

Ledum glandulosum Nutt. var. *glandulosum*

Leglg

Labrador tea

FORM: Erect shrub, 5-20dm (2-7ft) tall.

STEMS: Young branches finely hairy and dotted with glands, older branches smooth.

LEAVES: Alternate, evergreen, entire, ovate to elliptic, 2-6cm (1-2in) long, dark green above and light green and dotted with tiny yellow glands on lower surface, margins may be inrolled.

FLOWERS: Small, white or yellowish, in clusters at ends of branches; June-Aug.

FRUIT: Globose capsule, 1-3mm long.

SIMILAR SPECIES: *Menziesia ferruginea*, p.50, sticky, malodorous foliage; and *Rhododendron albiflorum*, p.59, no odor.

HABITAT: Riparian, bogs, and moist coniferous woods; indicator of wet, usually very acidic and nutrient poor organic soils.

NOTES: Fragrant when crushed.