

| | | | |
|-----------|-----------|--------|---|
| Pipo | Syal h.t. | 6 | |
| | Phma h.t. | 10 | |
| Psme | Caru h.t. | 10 | |
| | Aruv | 6 | |
| | Phma | 8 | |
| | Smst h.t. | 9 | |
| Abgr | Phma h.t. | Phma 7 | |
| | | Cooc 7 | |
| | Libo h.t. | Libo 5 | |
| | | Xete 7 | |
| | Asca h.t. | Asca 5 | |
| | | Tabr 2 | |
| | | Mefe 1 | |
| | Setr h.t. | 2 | |
| | Clun h.t. | Phma | 8 |
| | | Clun | 6 |
| | | Tabr | 3 |
| | | Mefe | 2 |
| Xete | | 7 | |
| Vagl | | 8 | |
| Xete h.t. | Cooc 7 | | |
| Thpl | Clun h.t. | Clun 5 | |
| | | Tabr 4 | |
| | | Mefe 3 | |
| | | Xete 3 | |
| | Asca h.t. | Asca 4 | |
| | | Tabr | |
| | | Mefe 3 | |
| | Gydr h.t. | 2 | |
| | Adpe h.t. | 2 | |
| | Atfi h.t. | Atfi 1 | |
| Opho h.t. | | | |
| Tshe | Clun h.t. | Clun 3 | |
| | | Arnu 4 | |
| | | Mefe 3 | |
| | | Xete 5 | |
| | Asca h.t. | Asca 3 | |
| | Arnu 3 | | |
| | Mefe | | |
| Gydr h.t. | + | | |
| Tsme | Clun h.t. | Mefe 1 | |
| | | Xete 2 | |
| | Mefe h.t. | Xete | |
| | | Luhi | |
| | Stam h.t. | Mefe 2 | |
| | Luhi | | |
| Xete h.t. | Vagl 3 | | |
| | Luhi | | |
| Abta | Clun h.t. | Clun 3 | |
| | | Mefe 1 | |
| | | Xete 5 | |
| | Mefe h.t. | Cooc | 1 |
| | | Xete | |
| | | Vasc | |
| | Stam h.t. | Luhi | |
| | | Mefe | |
| | Caca h.t. | Lica | 1 |
| | | Caca | 2 |
| Vaca | | | |
| Xete h.t. | Legl | 2 | |
| | Cooc | 5 | |
| | Vagl | 6 | |
| | Vasc | 4 | |
| | Luhi | | |
| Pico | Vasc h.t. | 9 | |
| | Xete c.t. | | |
| | Vaca c.t. | | |



ROSACEAE
Spiraea betulifolia var. *lucida* (Douglas.) Hitchc.
 Spbel

spiraea, shiny-leaf spiraea

FORM: Strongly rhizomatous, glabrous, medium shrub, 20-60cm (8-24in) tall.
 STEMS: Erect, slender, light yellow-brown and smooth, not hairy; aging to reddish-brown.
 LEAVES: Alternate, deciduous, ovate, 2-7cm (1-3in) long, generally coarsely toothed above middle, dark green on upper surface, slightly pale below, side veins of leaf bend toward apex.
 FLOWERS: Dull white in flat-topped corymbs, 3-8cm (1-3in) broad; June-July.
 FRUIT: Glabrous follicles, 3mm long (1).
 SIMILAR SPECIES: *Spiraea* species; or young *Amelanchier alnifolia*, p.36, different leaf serration and venation patterns.
 NOTES: Most common *Spiraea* species in northern Idaho.